

## QIBA Musculoskeletal (MSK) Biomarker Committee (BC) Call

Tuesday, February 18, 2019 at 10:30 AM CT

### Call Summary

#### In attendance

Xiaojuan Li, PhD (Co-Chair)

Thomas Link, MD, PhD (Co-Chair)

Michael Boss, PhD

Angie Botto-van Bemden, PhD

Robert Boutin, MD

Majid Chalian, MD

Alejandro Espinoza, PhD

Gabby Joseph, PhD

Feliks Kogan, PhD

Leon Lenchik, MD

Kecheng Liu, PhD, MBA

Nancy Obuchowski, PhD

Yuxi Pang, PhD

Erika Schneider, PhD

Ramya Srinivasan, MD

Carl Winalski, MD

Cory Wyatt, PhD

#### RSNA

Joe Koudelik

Susan Stanfa

**Moderator:** Dr. Link

#### **Presentation: Defining Normative MRI cartilage T<sub>2</sub> values in the knee: data from the Osteoarthritis Initiative (Dr. Gabby Joseph, PhD and Thomas Link, MD, PhD)**

[Some information taken from presentation slides]

- Two previous research studies were highlighted: T<sub>2</sub> Reference Database → T<sub>2</sub> Z-score quantification
- Joseph GB, McCulloch CE, Nevitt MC, Heilmeier U, Nardo L, Lynch JA, et al. [A reference database of cartilage 3 T MRI T<sub>2</sub> values in knees without diagnostic evidence of cartilage degeneration: data from the osteoarthritis initiative](#). *Osteoarthritis Cartilage*. 2015; 23(6):897-905. doi: 10.1016/j.joca.2015.02.006.
  - Purpose:
    - To establish a gender- and BMI-specific reference database of cartilage T<sub>2</sub> values
    - To assess the associations between cartilage T<sub>2</sub> values and gender, age, and BMI in knees without radiographic osteoarthritis or MRI-based (WORMS 0/1) evidence of cartilage degeneration
  - Why normative values?
    - Provide a reference database of T<sub>2</sub> values for future studies with similar acquisition and scanning methodologies
    - Enable comparisons to studies that focus on cohorts with expected abnormal T<sub>2</sub> values
  - Inclusion and exclusion criteria were provided and imaging methods and statistics were outlined
  - Results were provided:
    - Percentiles by gender and BMI
    - Adjusted T<sub>2</sub> values in males and females
    - T<sub>2</sub> and BMI in normal, overweight and obese subjects
    - The association between BMI and cartilage T<sub>2</sub> – figure showed adjusted means with 95% confidence intervals
    - The association between age and cartilage T<sub>2</sub> – figure showed adjusted means with 95% confidence intervals
  - Discussion:
    - Substantial variation found among subject and between cartilage compartments
    - While an association between cartilage T<sub>2</sub> and both age and gender was established, the association with BMI was the most pronounced (e.g., the higher BMI resulted in higher image noise)
    - The age and BMI relationship was not studied
    - Patients that were obese were more susceptible to developing cartilage defects and damage

- Age had very little impact on T<sub>2</sub> measurements; it was only significant in the patella and was borderline in the medial femur
  - There was not much difference in results between males and females
  - Interactions between muscle and cartilage were not addressed in this study
  - T<sub>2</sub> results were available only for the right knee
  - Ultimately, this study has data that are useful for only a specific technique – T<sub>2</sub>; data are not available for T1ρ
  
- Joseph GB, McCulloch CE, Nevitt MC, Gersing AS, Schwaiger BJ, Kretschmar M, et al. [Medial femur T<sub>2</sub> Z-scores predict the probability of knee structural worsening over 4-8 years: Data from the osteoarthritis initiative](#). *Journal of Magnetic Resonance Imaging*. 2017; 46(4):1128-1136. doi: 10.1002/jmri.25662.
  - Subject selection was detailed and methods were explained
  - Results regarding the association between baseline cartilage T<sub>2</sub> Z-score and the probability of progression of OA were provided; the higher the Z-score, the higher probability of cartilage degeneration in the patient's future
  - Discussion: knowing a T<sub>2</sub> Z-score may be beneficial for clinicians to identify patients at risk for OA progression; these patients would benefit most from modifiable lifestyle changes that slow OA progression
  - Factors that affect T<sub>2</sub> quantification:
    - Reference database/Z-score calculation feasible with the OAI data since the following factors were standardized:
      - MRI scanner
      - MRI pulse sequence
      - Field strength
      - T<sub>2</sub> fitting method
      - RF coil
  - Comment re: interest in data for joint space narrowing
  - Baseline to discuss data interpretation and how to deal with it; a good reference database exists for T<sub>2</sub>, but not for newer sequences and T1ρ
  - Suggestion to define Z-scores instead of absolute values
  - For standardization, a Z-score that is site-dependent rather than scanner dependent was suggested, since the absolute values between scanners show high variation
  - Caution voiced regarding having an imaging protocol/Profile be scanner dependent
  - Suggestion to use OA phantom
    - Caution to avoid overcorrecting data is needed
    - Both OAI and MAPSS sequences were used in the Arthritis Foundation (AF) phantom study
      - Systematic differences were identified; correlation and differences were seen between the two
      - The phantom used in the AF study was not an OAI phantom, but another calibration phantom
    - There was concern re: the life span of phantoms
    - Material components were solutions (not gel-based) and should be easy to reproduce
    - While solutions are typically quite stable; better reproducibility when using agar
  - **It was advised that the MSK Profile and protocol be site and scanner independent**
  - Z-scores are not as dependent on absolute values; data needed for comparison between sites

- Z-scores will be better than absolute values in regard to reducing variability and dependence from different scanners
- This discussion will inform the work on Section 3.9: Data Interpretation of the MSK Profile and will be continued during an upcoming MSK BC call

**Next Call:** Tuesday, March 26, 2019 at 10 AM CT

---

RSNA Staff attempt to identify and capture all committee members participating on WebEx calls. However, **if multiple callers join simultaneously or call in without logging on to the WebEx, identification is not possible.** Call participants are welcome to contact RSNA staff at [QIBA@RSNA.org](mailto:QIBA@RSNA.org) if their attendance is not reflected on the call summaries.