Test-Retest, Reproducibility, and/or Repeatability in Liver MRE Publications

- 1. Hines, C.D.G., et al., *Repeatability of magnetic resonance elastography for quantification of hepatic stiffness.* Journal of Magnetic Resonance Imaging, 2010. **31**: p. 725-731.
- 2. Hines, C.D.G., et al., *Effects of postprandial state and mesenteric blood flow on the repeatability of MR elastography in asymptomatic subjects.* Journal of Magnetic Resonance Imaging, 2011. **33**: p. 239-244.
- 3. Shire, N.J., et al., *Test-retest repeatability of MR elastography for noninvasive liver fibrosis assessment in Hepatitis C.* Journal of Magnetic Resonance Imaging, 2011. **34**: p. 947-955.
- 4. Mannelli, L., et al., *Magnetic resonance elastography: feasibility of liver stiffness measurements in healthy volunteers at 3T.* Clinical Radiology, 2012. **67**: p. 258-262.
- 5. Rustogi, R., et al., *Accuracy of MR elastography and anatomic MR imaging features in the diagnosis of severe hepatic fibrosis and cirrhosis.* Journal of Magnetic Resonance Imaging, 2012. **35**: p. 1356-1364.
- 6. Bohte, A.E., et al., *MR elastography of the liver: defining thresholds for detecting viscoelastic changes.* Radiology, 2013. **269**(3): p. 768-776.
- Lee, D.H., et al., *MR elastography of healthy liver parenchyma: normal value and reliability of the liver stiffness value measurement.* Journal of Magnetic Resonance Imaging, 2013. 38: p. 1215-1223.
- 8. Jajamovich, G.H., et al., *Quantitative liver MRI combining phase contrast imaging, elastography, and DWI: assessment of reproducibility and postprandial effect at 3.0 T.* PLoS ONE, 2014. **9**(5): p. e97355.
- 9. Lee, Y.J., et al., *MR elastography for noninvasive assessment of hepatic fibrosis: reproducibility of the examination and reproducibility and repeatability of the liver stiffness value measurement.* Journal of Magnetic Resonance Imaging, 2014. **39**: p. 326-331.
- 10. Serai, S., et al., *Cross-vendor validation of liver magnetic resonance elastography*. Abdominal Imaging, 2015. **40**: p. 789-794.
- 11. Shi, Y., et al., *Short- and midterm repeatability of magnetic resonance elastography in healthy volunteers at 3.0 T.* Magnetic Resonance Imaging, 2014. **32**: p. 665-670.
- Shin, S.U., et al., Prediction of esophageal varices in patients with cirrhosis: usefulness of three- dimensional MR elastography with echo-planar imaging technique. Radiology, 2014.
  272(1): p. 143-153.
- 13. Shinagawa, Y., et al., *Optimization of scanning parameters for MR elastography at 3.0T clinical unit: volunteer study.* Jpn J Radiol, 2014. **32**: p. 441-446.
- 14. Venkatesh, S.K., et al., Magnetic resonance elastography of liver in healthy asians: normal liver stiffness quantification and reproducibility assessment. Journal of Magnetic Resonance Imaging, 2014. 39: p. 1-8